Neuroscience Letters, 119 (1990) 277–280
Elsevier Scientific Publishers Ireland Ltd.

AUTHOR INDEX

Abreu, P., see Moujir, F., 12
Agerberth, B., see Aguirre, J.A., 129
Agnati, L.F., see Aguirre, J.A., 129
Ago, Yu., see Ishikawa, T., 114
Aguirre, J.A., Fuxe, K., Agnati, L.F., Agerberth, B. and Mutt, V., Vasopressor effects of intracerebroventricular injections of PEC-60, a novel porcine 60-residue intestinal peptide, in the awake unrestrained male rat, 129
Akaike, N., see Oyama, Ya., 261
Alonso, R., see Moujir, F., 12
Ansselin, A.D., Westland, K. and Pollard, J.D., Low dose, short term Cyclosporin A does not protect the Schwann cells of allogeneic nerve grafts, 219
Aoki, E., see Seto-Ohshima, A., 60
Arai, A., Kessler, M. and Lynch, G., The effects of adenosine on the development of long-term potentiation, 41
Atlas, D., see Weiss, C., 241
Azuma, H., Yamatodani, A., Yagi, A., Nishimura, T. and Wada, H., Influence of intensity of illumination during the light period on diurnal variations of pineal indoles in rats and mice, 15

Barron, E., see Perlmutter, L.S., 32
Basbaum, A.I., see Levine, J.D., 37
Beckley, D.J., see Singh, J., 27
Benecke, R., see Dressler, D., 153
Beracochea, D., see Villalobos, J., 97
Berría, M.I., see Castagnino, C.G., 132
Bigbee, J.W., Yu, D.S. and Yu, R.K., Morphometric analysis of the developing optic nerve of the F1 heterotic mouse and its parental strains, 179
Bingren, H., see Cardell, M., 228
Blitzer, R.D., Gil, O. and Landau, E.M., Cholinergic stimulation enhances long-term potentiation in the CA1 region of rat hippocampus, 207
Bobillier, P., see Sanne, J. -L., 94
Boer, G.J., see Van Veldhuizen, M.J.A., 233
Bordón, R., see Moujir, F., 12
Bout, R.G., see Dubbeldam, J.L., 223
Bugiani, O., Giaccone, G., Verga, L., Pollo, B., Ghetti, B., Frangione, B. and Tagliavini, Fabrizio, Alzheimer patients and Down patients: abnormal presynaptic terminals are related to cerebral preamyloid deposits, 56
Burnstock, G., see Crowe, R., 203

Cardell, M., Bingren, H., Wieloch, T., Zivin, J. and Saitoh, T., Protein kinase C is translocated to cell membranes during cerebral ischemia, 228
Castagnino, C.G. and Berría, M.I., Fibronectin involvement following herpes simplex infection of cultured rat astrocytes, 132
Ceci, A., see French, E.D., 159
Cervera, P., see Hauw, J. -J., 182
Chui, H.C., see Perlmutter, L.S., 32
Clayworth, C., see Singh, J., 27
Coderre, T.J., see Levine, J.D., 37
Crowe, R. and Burnstock, G., The subserosal ganglia of the human taenia, 203
.D. Lund, R., see .E. Westrum, L., 265

Darnall, R.A. and Guyenet, P., Respiratory modulation of pre- and postganglionic lumbar vasomotor sympathetic neurons in the rat, 148
Delaère, P., see Hauw, J. -J., 182
Denavit-Saubié, M., see Pierrefiche, O., 90
Doll, C.J., see Reiner, P.B., 175
Dorman, R.V., see Terrian, D.M., 211
Dorsa, D.M., see Szot, P., 215
Dreier, J.P. and Heinemann, U., Late low magnesium-induced epileptiform activity in rat entorhinal cortex slices becomes insensitive to the anticonvulsant valproic acid, 68
Dressler, D., Voth, E., Feldmann, M. and Benecke, R., Safety aspects of transcranial brain stimulation in man tested by single photon emission-computed tomography, 153
Dubbeldam, J.L. and Bout, R.G., The identification of the motor nuclei innervating the tongue muscles in the mallard (*Anas platyrhynchos*); an HRP study, 223
Dutton, G.R., see Tigges, G.A., 23
Duyckaerts, C., see Hauw, J. -J., 182
.E. Westrum, L., J.N. Kott, , H. Vickland, , M.H. Hankin, and .D. Lund, R., Fetal olfactory bulb transplants send projections to host olfactory cortex in the rat, 265

Ehrhart-Bornstein, M., Thorn, N.A. and Treiman, M., Chronic osmotic stimulation reduces vasopressin but not synaptophysin content in rat neurohypophysis, 122
Eikelenboom, P., see Rozemuller, J.M., 75
Elazar, Z., Ganchrow, D., Paz, M., Rabinowitz, R., Paz, Z. and Korczyn, A.D., Dopaminergic drugs influence the intensity of catalepsy induced by microinjections of carbachol into the reticular formation, 245
El-Beheiry, H. and Puil, E., Unusual features of GABA responses in layers IV–V neurons of neocortex, 83
Emson, P.C., see Seto-Ohshima, A., 60

Feenstra, M.G.P., see Van Veldhuizen, M.J.A., 233
Feldmann, M., see Dressler, D., 153
Ferris, C.F., see Szot, P., 215
Finkbeiner, W.E., see Levine, J.D., 37
Fishman, P.H., see Gordon, E.A., 187
Fossom, L.H., see Gupta, M., 49
Foutz, A.S., see Pierrefiche, O., 90
Fox, L.M., see McNulty, J.A., 237
Frangione, B., see Bugiani, O., 56
Franke, C., see Quasthoff, S., 191
French, E.D. and Ceci, A., Non-competitive *N*-methyl-D-aspartate antagonists are potent activators of ventral tegmental A_{10} dopamine neurons, 159
Fujimori, K., see Nakazawa, K., 5
Fujimoto, K., see Kawashima, K., 156
Fuxe, K., see Aguirre, J.A., 129

Ganchrow, D., see Elazar, Z., 245
Gannon, R.L., see Terrian, D.M., 211
Gean, P.-W., NMDA receptor-independent epileptiform activity induced by magnesium-free solution in rat amygdala neurons is blocked by CNQX, 53
Ghetti, B., see Bugiani, O., 56
Giaccone, G., see Bugiani, O., 56
Gil, O., see Blitzer, R.D., 207

Gordon, E.A, Krstenansky, J.L. and Fishman, P.H., Centrally truncated neuropeptide Y analog acts as an agonist for Y_1 receptors on SK-N-MC cells, 187
Grange, E., see Sanne, J. -L., 94
Gündel, J., Steinhäuser, C. and Matthies, H., Proteolytic enzymes do not destroy the *N*-methyl-D-aspartate (NMDA) sensitivity of acutely isolated hippocampal CA1 and CA3 neurons from postnatal rats, 249
Gupta, M., Stoler, M.H., Fossom, L.H. and Tank, A.W., Tyrosine hydroxylase mRNA in the dopaminergic neurons of young adult and aged mice by in situ hybridization, 49
Guyenet, P., see Darnall, R.A., 148
H. Vickland, , see .E. Westrum, L., 265

Haroian, A.J., see Rubertone, J.A., 79
Harrison, N.L., Lovinger, D.M., Lambert, N.A., Teyler, T.J., Prager, R., Ong, J. and Kerr, D.I.B., The actions of 2-hydroxy-saclofen at presynaptic $GABA_B$ receptors in the rat hippocampus, 272
Hasegawa, T. and Sakuma, Ya., Estrogen-induced suppression of female rat forebrain neurons with axons to ventral midbrain, 171
Hatta, K., see Hori, T., 9
Hatt, H., see Quasthoff, S., 191
Hauw, J. -J., Verny, M., Delaère, P., Cervera, P., He, Y. and Duyckaerts, C., Constant neurofibrillary changes in the neocortex in progressive supranuclear palsy. Basic differences with Alzheimer's disease and aging, 182
Heinemann, U., see Dreier, J.P., 68
Heizmann, C.W., see Seto-Ohshima, A., 60
Hernandez, G., see Moujir, F., 12
He, Y., see Hauw, J. -J., 182
Higuchi, T., see Honda, K., 167
Hirano, T., Depression and potentiation of the synaptic transmission between a granule cell and a Purkinje cell in rat cerebellar culture, 141
Hirano, T., Effects of postsynaptic depolarization in the induction of synaptic depression between a granule cell and a Purkinje cell in rat cerebellar culture, 145
Hiranuma, K., see Isu, N., 163
Hökfelt, T., see Meister, B., 136
Honda, K., Negoro, H., Higuchi, T. and Takano, S., Activation of supraoptic neurosecretory cells by local osmotic stimulation of the median preoptic nucleus, 167
Hori, T., Yamamoto, T., Hatta, K. and Moroji, T., Biphasic effects of magnesium on the [^{3}H]N-(1-(2-thienyl)cyclohexyl)- 3,4-piperidine binding in the rat cerebral cortex, 9
Hugon, J. and Vallat, J.M., Abnormal distribution of phosphorylated neurofilaments in neuronal degeneration induced by kainic acid, 45

Ichikawa, T., see Isu, N., 163
Imamura, K., Mori, K., Oka, S. and Watanabe, Ya., Variations by layers and developmental changes in expression of telencephalin in the visual cortex of cat, 118
Inoue, K., see Nakazawa, K., 5
Ishikawa, T., Nagata, S., Ago, Yu., Takahashi, K. and Karibe, M., The central inhibitory effect of interleukin-1 on gastric acid secretion, 114
Ishizaki, Ya., see Kawashima, K., 156
Isu, N., Sakuma, A., Hiranuma, K., Ichikawa, T. and Uchino, Y., Localization and synaptic effects of inhibitory vestibulocollic neurons activated by the posterior semicircular canal nerve in the cat, 163

Jaffard, R., see Villalobos, J., 97
J.N. Kott, , see .E. Westrum, L., 265
Jürgens, H., see Valen, F.V., 195

Karibe, M., see Ishikawa, T., 114
Kawashima, K., Watanabe, N., Oohata, H., Fujimoto, K., Suzuki, T., Ishizaki, Ya., Morita, I. and Murota, S., Synthesis and release of acetylcholine by cultured bovine arterial endothelial cells, 156
Kerr, D.I.B., see Harrison, N.L., 272
Kessler, M., see Arai, A., 41
Khateb, A., see Serafin, M., 125
Kimura, A., Sato, A. and Takano, Yu., Stimulation of the nucleus basalis of Meynert does not influence glucose utilization of the cerebral cortex in anesthetized rats, 101
Knight, R.T., see Singh, J., 27
Koistinaho, J., see Roivainen, R., 105
Korczyn, A.D., see Elazar, Z., 245
Krekel, J., Weihe, E., Nohr, D., Yanaihara, N. and Weber, E., Distribution of Met-enkephalyl-Arg-Gly-Leu in rat larynx: partial coexistence with vasoactive intestinal polypeptide, peptide histidine isoleucine and neuropeptide Y, 64
Krstenansky, J.L., see Gordon, E.A., 187
Kudo, Y., see Yuzaki, M., 19

Lambert, N.A., see Harrison, N.L., 272
Landau, E.M., see Blitzer, R.D., 207
Laycock, A.G., see Reiner, P.B., 175
Levine, J.D., Coderre, T.J., White, D.M., Finkbeiner, W.E. and Basbaum, A.I., Denervation-induced inflammation in the rat, 37
Li, Yu.-Q., Rao, Z.-R. and Shi, J.-W., Midbrain periaqueductal gray neurons with substance P- or enkephalin-like immunoreactivity send projection fibers to the nucleus accumbens in the rat, 269
Lovinger, D.M., see Harrison, N.L., 272
Lynch, G., see Arai, A., 41

Matthies, H., see Gündel, J., 249
McNulty, J.A., Fox, L.M. and Spurrier, W.A., A circannual cycle in pinealocyte synaptic ribbons in the hibernating and seasonally reproductive 13-lined ground squirrel (*Spermophilus tridecemlineatus*), 237
Mehler, W.R., see Rubertone, J.A., 79
Meister, B., Scanlon, M.F. and Hökfelt, T., Occurrence of galanin-like immunoreactivity in growth hormone-releasing factor (GRF)-containing neurons of the monkey *(Macaca fascicularis)* infundibular nucleus and median eminence, 136
M.H. Hankin, , see .E. Westrum, L., 265
Mikoshiba, K., see Yuzaki, M., 19
Mitchell, D., see Sher, GrahamD., 199
Miyawaki, K.A.A., see Yuzaki, M., 19
Mori, K., see Imamura, K., 118
Morita, I., see Kawashima, K., 156
Moroji, T., see Hori, T., 9
Moujir, F., Bordón, R., Santana, C., Abreu, P., Hernandez, G. and Alonso, R., Ovarian steroids block the isoproterenol-induced elevation of pineal melatonin production in the female rat, 12
Mühlethaler, M., see Serafin, M., 125
Murota, S., see Kawashima, K., 156
Mutt, V., see Aguirre, J.A., 129

Nagata, S., see Ishikawa, T., 114
Nagatsu, T., see Yoshida, M., 109
Nakai, Ya., see Takeda, F., 86
Nakazawa, K., Inoue, K., Fujimori, K. and Takanaka, A., ATP-activated single-channel currents recorded from cell-free patches of

pheochromocytoma PC12 cells, 5
Necker, R., Cells of origin of ascending and descending as well as branching fibers in the cervical spinal cord of the pigeon, 1
Negoro, H., see Honda, K., 167
Nishimura, T., see Azuma, H., 15
Niwa, T., see Yoshida, M., 109
Noguchi, K., see Zhang, J.-H., 257
Nohr, D., see Krekel, J., 64

Oka, S., see Imamura, K., 118
Ong, J., see Harrison, N.L., 272
Oohata, H., see Kawashima, K., 156
Otte-Höller, I., see Stekhoven, J.H.S., 71
Oyama, Ya. and Akaike, N., Triphenyltin: a potent excitatory neurotoxicant. Its reciprocal effects on voltage-dependent Na and K currents of mammalian brain neuron, 261

Paz, M., see Elazar, Z., 245
Paz, Z., see Elazar, Z., 245
Peers, C., Hypoxic suppression of K currents in type I carotid body cells: selective effect on the Ca^2-activated K current, 253
Perlmutter, L.S., Barron, E. and Chui, H.C., Morphologic association between microglia and senile plaque amyloid in Alzheimer's disease, 32
Philibert, R.A., see Tigges, G.A., 23
Piechot, G., see Valen, F.V., 195
Pierrefiche, O., Foutz, A.S. and Denavit-Saubié, M., Pneumotaxic mechanisms in the non-human primate: effect of the *N*-methyl-D-aspartate (NMDA) antagonist ketamine, 90
Pollard, J.D., see Ansselin, A.D., 219
Pollo, B., see Bugiani, O., 56
Prager, R., see Harrison, N.L., 272
Puil, E., see El-Beheiry, H., 83

Quasthoff, S., Franke, C., Hatt, H. and Richter-Turtur, M., Two different types of potassium channels in human skeletal muscle activated by potassium channel openers, 191

Rabinowitz, R., see Elazar, Z., 245
Rao, Z.-R., see Li, Yu.-Q., 269
Reiner, P.B., Laycock, A.G. and Doll, C.J., A pharmacological model of ischemia in the hippocampal slice, 175
Renkawek, K., see Schuurmans Stekhoven, J.H.S., 71
Richter-Turtur, M., see Quasthoff, S., 191
Roivainen, R. and Koistinaho, J., Decentralization induces long-term c-fos protein-like immunoreactivity in non-neuronal cells in the rat superior cervical ganglion, 105
Rozemuller, J.M., Stam, F.C. and Eikelenboom, P., Acute phase proteins are present in amorphous plaques in the cerebral but not in the cerebellar cortex of patients with Alzheimer's disease, 75
Rubertone, J.A., Haroian, A.J., Vincent, S.L. and Mehler, W.R., The rat parvocellular reticular formation: I. Afferents from the cerebellar nuclei, 79

Saitoh, T., see Cardell, M., 228
Sakuma, A., see Isu, N., 163
Sakuma, Ya., see Hasegawa, T., 171
Sanne, J. -L., Scarna, H., Grange, E. and Bobillier, P., Restraint stress and adrenalectomy do not affect the level of rat cerebral enolase, 94
Santana, C., see Moujir, F., 12
Sato, A., see Kimura, A., 101
Sato, M., see Zhang, J.-H., 257
Sato, T., see Takeda, F., 86
Scanlon, M.F., see Meister, B., 136
Scarna, H., see Sanne, J. -L., 94
Schuurmans Stekhoven, J.H.S., Renkawek, K., Otte-Höller, I. and Stols, A., Exogenous aluminum accumulates in the lysosomes of cultured rat cortical neurons, 71
Sela, D., see Weiss, C., 241
Semba, R., see Seto-Ohshima, A., 60
Serafin, M., Khateb, A. and Mühlethaler, M., Opiates inhibit pedunculopontine neurones in guinea pig brainstem slices, 125
Seto-Ohshima, A., Aoki, E., Semba, R., Emson, P.C. and Heizmann, C.W., Parvalbumin immunoreactivity in the central auditory system of the gerbil: a developmental study, 60
Sher, G.D. and Mitchell, D., Intrathecal *N*-methyl-D-aspartate induces hyperexcitability in rat dorsal horn convergent neurones, 199
Shi, J.-W., see Li, Yu.-Q., 269
Shioda, S., see Takeda, F., 86
Singh, J., Knight, R.T., Woods, D.L., Beckley, D.J. and Clayworth, C., Lack of age effects on human brain potentials preceding voluntary movements, 27
Spurrier, W.A., see McNulty, J.A., 237
Stam, F.C., see Rozemuller, J.M., 75
Steinhäuser, C., see Gündel, J., 249
Stoler, M.H., see Gupta, M., 49
Stols, A., see Stekhoven, J.H.S., 71
Suzuki, T., see Kawashima, K., 156
Szot, P., Ferris, C.F. and Dorsa, D.M., [^{3}H]Arginine-vasopressin binding sites in the CNS of the golden hamster, 215

Tagliavini, Fabrizio, see Bugiani, O., 56
Takahashi, K., see Ishikawa, T., 114
Takanaka, A., see Nakazawa, K., 5
Takano, S., see Honda, K., 167
Takano, Yu., see Kimura, A., 101
Takeda, F., Takeda, M., Sato, T., Shioda, S. and Nakai, Ya., Distribution and co-existence of Met-enkephalin-like and mesotocin-like immunoreactivity in the neural lobe of the pituitary of the frog, 86
Takeda, M., see Takeda, F., 86
Tank, A.W., see Gupta, M., 49
Terrian, D.M., Dorman, R.V. and Gannon, R.L., Characterization of the presynaptic calcium channels involved in glutamate exocytosis from rat hippocampal mossy fiber synaptosomes, 211
Teyler, T.J., see Harrison, N.L., 272
Thorn, N.A., see Ehrhart-Bornstein, M., 122
Tigges, G.A., Philibert, R.A. and Dutton, G.R., K- and temperature-evoked taurine efflux from hypothalamic astrocytes, 23
Tohyama, M., see Zhang, J.-H., 257
Treiman, M., see Ehrhart-Bornstein, M., 122

Uchino, Y., see Isu, N., 163

Valen, F.V., Piechot, G. and Jürgens, H., Calcitonin gene-related peptide (CGRP) receptors are linked to cyclic adenosine monophosphate production in SK-N-MC human neuroblastoma cells, 195
Vallat, J.M., see Hugon, J., 45
Van Veldhuizen, M.J.A., Feenstra, M.G.P., Boer, G.J. and Westerink, B.H.C., Microdialysis studies on cortical noradrenaline release: basic characteristics, significance of extracellular calcium and massive post-mortem increase, 233
Verga, L., see Bugiani, O., 56
Verny, M., see Hauw, J. -J., 182
Villalobos, J., Beracochea, D. and Jaffard, R., Age-related changes in the subcortical afferents to the medial frontal cortex in mice: a WGA–HRP study, 97

Vincent, S.L., see Rubertone, J.A., 79
Voth, E., see Dressler, D., 153

Wada, H., see Azuma, H., 15
Watanabe, N., see Kawashima, K., 156
Watanabe, Ya., see Imamura, K., 118
Weber, E., see Krekel, J., 64
Weihe, E., see Krekel, J., 64
Weiss, C., Sela, D. and Atlas, D., Divalent cations effectively replace Ca^{2+} and support bradykinin induced noradrenaline release, 241
Westerink, B.H.C., see Van Veldhuizen, M.J.A., 233
Westland, K., see Ansselin, A.D., 219
White, D.M., see Levine, J.D., 37
Wieloch, T., see Cardell, M., 228
Woods, D.L., see Singh, J., 27

Yagi, A., see Azuma, H., 15
Yamamoto, T., see Hori, T., 9
Yamatodani, A., see Azuma, H., 15
Yanaihara, N., see Krekel, J., 64
Yoshida, M., Niwa, T. and Nagatsu, T., Parkinsonism in monkeys produced by chronic administration of an endogenous substance of the brain, tetrahydroisoquinoline: the behavioral and biochemical changes, 109
Yu, D.S., see Bigbee, J.W., 179
Yu, R.K., see Bigbee, J.W., 179
Yuzaki, M., Kudo, Y., Miyawaki, K.A.A. and Mikoshiba, K., MK-801 blocked the functional NMDA receptors in identified cerebellar neurons, 19

Zhang, J.-H., Sato, M., Noguchi, K. and Tohyama, M., The differential expression patterns of the mRNAs encoding β subunits (β_1, β_2 and β_3) of $GABA_A$ receptor in the olfactory bulb and its related areas in the rat brain, 257
Zivin, J., see Cardell, M., 228

CONTENTS *(continued from back cover)*

A circannual cycle in pinealocyte synaptic ribbons in the hibernating and seasonally reproductive 13-lined ground squirrel (*Spermophilus tridecemlineatus*)
J.A. McNulty, L.M. Fox and W.A. Spurrier (U.S.A.) 237
Divalent cations effectively replace Ca^{2+} and support bradykinin induced noradrenaline release
C. Weiss, D. Sela and D. Atlas (Israel) 241
Dopaminergic drugs influence the intensity of catalepsy induced by microinjections of carbachol into the reticular formation
Z. Elazar, D. Ganchrow, M. Paz, R. Rabinowitz, Z. Paz and A.D. Korczyn (Israel) 245
Proteolytic enzymes do not destroy the *N*-methyl-D-aspartate (NMDA) sensitivity of acutely isolated hippocampal CA1 and CA3 neurons from postnatal rats
J. Gündel, C. Steinhäuser and H. Matthies (G.D.R.) 249
Hypoxic suppression of K currents in type I carotid body cells: selective effect on the Ca^2-activated K current
C. Peers (U.K.) 253
The differential expression patterns of the mRNAs encoding β subunits (β_1, β_2 and β_3) of $GABA_A$ receptor in the olfactory bulb and its related areas in the rat brain
J.-H. Zhang, M. Sato, K. Noguchi and M. Tohyama (Japan) 257
Triphenyltin: a potent excitatory neurotoxicant. Its reciprocal effects on voltage-dependent Na and K currents of mammalian brain neuron
Ya. Oyama and N. Akaike (Japan) 261
Fetal olfactory bulb transplants send projections to host olfactory cortex in the rat
L. .E. Westrum, J.N. Kott, H. Vickland, M.H. Hankin and R. .D. Lund (U.S.A.) 265
Midbrain periaqueductal gray neurons with substance P- or enkephalin-like immunoreactivity send projection fibers to the nucleus accumbens in the rat
Yu.-Q. Li, Z.-R. Rao and J.-W. Shi (People's Republic of China) 269
The actions of 2-hydroxy-saclofen at presynaptic $GABA_B$ receptors in the rat hippocampus
N.L. Harrison, D.M. Lovinger, N.A. Lambert, T.J. Teyler, R. Prager, J. Ong and D.I.B. Kerr (U.S.A., Australia) 272

Author Index 277